



Better soils. Better crops. Better stock.™

BioAg RumiMate®

Nutritional supplement for ruminants and other animals

Product overview

BioAg liquid cultures are a revolutionary way of delivering essential nutrients and metabolites to your livestock, as well as stimulating microbial activity in the rumen and stomach. The end result is enhanced digestion.

BioAg RumiMate is formulated to stimulate microbial activity in the animal's digestive tract. It enhances the natural replenishment of digestive microflora that occurs through grazing. This fermented microbial culture contains a rich source of vitamins, minerals, proteins, enzymes, amino acids and carbohydrates.

BioAg RumiMate can be administered by oral drench, mixed with feed rations or mixed with drinking water. It is suitable for lead feeding stock before ration changes or continual use for intensive animals.

At a glance

- Balanced nutrient supply for livestock
- Stimulates digestive microbial activity
- Enhances natural digestive microflora
- Administer by oral drench or mix with drinking water or feed rations



BioAg RumiMate® – Directions for use

Animal	Situation	Dosage per head per day	
		Stress (up to first 4 days)	Maintenance (ongoing)
Cattle	Milking	20 mL	6–10 mL
	Beef & dry cows	15–20 mL	3–7 mL
	Calves	8–12 mL	3–4 mL
Sheep	Wethers	8–15 mL	4–5 mL
	Ewes	8–15 mL	4–5 mL
	Lambs	5–8 mL	2–4 mL
Horses	Stud or mares	16 mL	4–8 mL
	Fillies	10–12 mL	5–6 mL
	Colts	8 mL	2–4 mL
Pigs	Sows	20 mL	6–10 mL
	Piglets	3–4 mL	1–2 mL
	Weaners	6 mL	2–3 mL
	Growers	10–15 mL	4–7 mL

The dosage provided will depend on a range of factors including the animal's weight physiological state and health; feed availability and quality (e.g. pasture, grain, hay, silage); and environmental conditions.

Use stress rate when stock are not performing normally and revert to maintenance rate after two days.

Mixing and administration

- Agitate contents well before use
- Refer to table above for application rate
- Apply by ONE of the following methods:
 - oral drench: use undiluted product and standard drenching procedures
 - mix with drinking water: add directly to water troughs, ensuring stock receive correct dosage based on standard drinking volume
 - mix with feed ration: dilute with water and mix with feed ration at total volume of 5 litres per tonne of feed.
 - Dilute product with water before mixing with feed ration
 - Use diluted product within 48 hours

Storage and handling

- Store all containers out of direct sunlight
- Shelf life: 6 months

Withholding Period:

- Nil



Better soils. Better crops. Better stock.™

**For more information,
phone 0800 224 624 or visit www.bioag.co.nz**

©Registered trademarks of BioAg Pty Ltd, ABN 58 086 880 211 ™Trademark of BioAg Pty Ltd, ABN 58 086 880 211 WORDSMITH27226